



Probing the Interstellar Dust of Galaxies at Redshift $\sim 0.4 - 0.8$ in the AstroSat UV Deep Field

Chayan Mondal

S. N. Bose National Centre for Basic Sciences, Kolkata

In collaboration with

Kanak Saha, Chian Chou Chen, Rogier A. Windhorst, Rolf A. Jansen, Thomas S.-Y. Lai, Irene Shivaiei, Hiddo S. B. Algera, and Bovornpratch Vijarnwannaluk

NUVA eMeeting 2026


8 October 2026



What is a 'Deep Field'?

A region of the dark sky that contains very few bright stars and have ample observing time which enables studying faint deep sky objects at different redshifts.


Taking pictures of 'Nothing'




The total amount of exposure time was
~272 hours (400 Hubble orbits)

*Credit: NASA, ESA, and S. Beckwith
(STScI) and the HUDF Team*



Redshift= 0.0



Why multi-band observation?






Redshift	wave_emitted	wave_observed
0.5	1500	2250 (UV)
4	1500	7500 (Optical)
10	1500	16500 (Near-IR)
15	1500	24000 (Far-IR)



AstroSat/UVIT – unique instrument to probe UV sky

Ultra-Violet Imaging Telescope (UVIT)

- Simultaneous imaging in FUV and NUV
- Multiple filters in FUV and NUV channels
- Angular resolution ~ 1.2 arcsec



AstroSat UV Deep Fields

2016

2022

2023

GOODS-South
(AUDFs)


NEP (JWST NEP
collaboration)

MACS J0416.1-2403

GOODS-North
(AUDFn)

AEGIS


Abell 2744
Abell S1063



Probing interstellar dust

Interstellar dust contributes only about 1% to the total ISM mass in galaxies



Though, **negligible in mass**, dust plays **outside role** in **shaping the galaxy SED** that we observe



Indirect way of probing dust content in galaxies at higher redshifts


Direct way of probing dust content in galaxies

Cosmic evolution of dust extinction



To probe UV continuum at $z \sim 0.6$, we need deep NUV imaging

Our measurement from 465 UV-selected galaxies at $z \sim 0.6$



Our measurements fill this gap that connects the cosmic noon to the present day. It further reinforces the gradual reddening of galaxy's UV continuum with cosmic time



Mondal et al. 2023b

Properties of small dust grains


Polycyclic Aromatic Hydrocarbon (PAH)

What are PAHs?

- Carbon-based smaller dust particles consisting of ~ 10 to a few hundred carbon atoms
- Contribute up to ~ 15% of the total carbon content in galaxies and up to ~5% of the total dust mass



PAH emission




MIR emission features:
Single photon stochastic
heating of smaller dust
particles (eg., PAH)

FIR continuum: Larger
grains in thermal
equilibrium with the
radiation field


PAH: Galaxy evolution

→ PAH fraction increases with gas-phase metallicity



Aniano+2020

→ PAH luminosity can trace SFR of galaxies




Kim+2024

PAH - Indirect probe of molecular gas


→ PAH intensity correlates well with molecular gas content

Nearby galaxies




Cortzen+2019

Distant galaxies




Shivaei+2024b

Current PAH studies & the upcoming PRIMA telescope





Our Objective


Q1. What drives
PAH enrichment in
galaxies at redshift
 $z \sim 0.5 - 0.8$?



Probing PAH using JWST MIRI at $z \sim 0.65$




Diagnostic parameter to probe PAH emission at $z \sim 0.65$




$$f_{\text{PAH}} = \frac{E}{0.5 \times (C1 + C2)}$$

$$f_{\text{PAH}} = \frac{\text{Flux}_{\text{F1280W}}}{0.5 \times (\text{Flux}_{\text{F770W}} + \text{Flux}_{\text{F2550W}})}$$



f_{PAH} (measured from MIRI fluxes) can probe the PAH mass fraction at $z \sim 0.65$


Relation with galaxy stellar mass



The PAH mass fraction of star-forming galaxies show a strong dependence on galaxy stellar mass

However, the massive quiescent galaxies show low PAH mass fraction

Relation with FUV luminosity





f_{PAH} increases with stellar mass and rest-frame FUV luminosity, identifying massive, UV-bright galaxies to have higher PAH mass fraction


Relation with ISM ionization

we observe an anti-correlation between f_{PAH} and the O32 emission line ratio, suggesting PAH photo-dissociation in strongly ionized ISM

Star-forming dwarf galaxies are more prone to host highly ionized ISM which is less favourable for PAH survival



Relation with gas-phase metallicity




PAH enrichment in galaxies goes through a rapid transition phase from a PAH-poor to a PAH-rich system, at about 70% solar metallicity

Low-mass metal-poor SFGs, which are increasingly common at even higher redshifts, host more extreme conditions less favorable for PAH production and survival

To Summarize

- For SFGs, galaxy stellar mass and UV luminosity together play the most important role in regulating PAH enrichment and emission
- However, SFGs with harder ISM radiation field can undergo efficient photo-dissociation of PAH molecules
- QGs show relatively lower PAH mass fraction, indicating that the quenching process that affected star formation in these systems could also destroy PAH molecules



More exploration ongoing in multiple AstroSat UV Deep Fields....

